



# AR $\alpha$ 2A rabbit pAb

Cat No.:ES4893

For research use only


## Overview

<b>Product Name</b>	AR $\alpha$ 2A rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse;Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Adrenergic Receptor alpha-2A. AA range:331-380
<b>Specificity</b>	AR $\alpha$ 2A Polyclonal Antibody detects endogenous levels of AR $\alpha$ 2A protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Alpha-2A adrenergic receptor
<b>Gene Name</b>	ADRA2A
<b>Cellular localization</b>	Cell membrane ; Multi-pass membrane protein .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	48kD
<b>Human Gene ID</b>	150
<b>Human Swiss-Prot Number</b>	P08913
<b>Alternative Names</b>	ADRA2A; ADRA2R; ADRAR; Alpha-2A adrenergic receptor; Alpha-2 adrenergic receptor subtype C10; Alpha-2A adrenoreceptor; Alpha-2A adrenoceptor; Alpha-2AAR
<b>Background</b>	Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. They include






3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. These receptors have a critical role in regulating neurotransmitter release from sympathetic nerves and from adrenergic neurons in the central nervous system. Studies in mouse revealed that both the alpha2A and alpha2C subtypes were required for normal presynaptic control of transmitter release from sympathetic nerves in the heart and from central noradrenergic neurons; the alpha2A subtype inhibited transmitter release at high stimulation frequencies, whereas the alpha2C subtype modulated neurotransmission at lower levels of nerve activity. This gene encodes alpha2A subtype and it contains no introns in either its coding or untranslated sequences. [provided by RefSeq, Jul 2008],





Western Blot analysis of Jurkat cells using AR  $\alpha$ 2A Polyclonal Antibody

Immunofluorescence analysis of NIH/3T3 cells, using Adrenergic Receptor alpha-2A Antibody. The picture on the right is blocked with the synthesized peptide.





Immunohistochemistry analysis of paraffin-embedded human testis tissue, using Adrenergic Receptor alpha-2A Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat and LOVO cells, using Adrenergic Receptor alpha-2A Antibody. The lane on the right is blocked with the synthesized peptide.

