

eIF4B rabbit pAb

Cat No.:ES5075

For research use only

Overview

Product Name eIF4B rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunofluoroscopco: 1/200 - 1/1000 EU

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human eIF4B. AA

range:388-437

Specificity eIF4B Polyclonal Antibody detects endogenous

levels of eIF4B protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Eukaryotic translation initiation factor 4B

Gene Name EIF4B

Cellular localization cytosol,polysome,eukaryotic translation initiation

factor 4F complex, dendrite, neuronal cell body,

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band70kDHuman Gene ID1975Human Swiss-Prot NumberP23588

Alternative Names EIF4B; Eukaryotic translation initiation factor 4B;

eIF-4B

Background function:Required for the binding of mRNA to

ribosomes. Functions in close association with EIF4-F and EIF4-A. Binds near the 5'-terminal cap of




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com





mRNA in presence of EIF-4F and ATP. Promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both EIF4-A and EIF4-F.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Self-associates and interacts with EIF3 p170 subunit.,

Immunofluorescence analysis of A549 cells, using eIF4B Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using eIF4B Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from HepG2 and HUVEC cells, using eIF4B Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com