

## EphA4 (phospho Tyr596) rabbit pAb

**Cat No.: ES5128** 

For research use only

## Overview

Product Name EphA4 (phospho Tyr596) rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen Synthesized phospho-peptide around the

phosphorylation site of human EphA4 (phospho

Tyr596)

**Specificity** Phospho-EphA4 (Y596) Polyclonal Antibody detects

endogenous levels of EphA4 protein only when

phosphorylated at Y596.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Ephrin type-A receptor 4

Gene Name EPHA4

**Cellular localization** Cell membrane ; Single-pass type I membrane

protein . Cell projection, axon . Cell projection, dendrite . Cell junction, synapse, postsynaptic density membrane . Early endosome . Cell junction, adherens junction . Clustered upon activation and

targeted to

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 110kD
Human Gene ID 2043
Human Swiss-Prot Number P54764

Alternative Names EPHA4; HEK8; SEK; TYRO1; Ephrin type-A receptor 4;

EPH-like kinase 8; EK8; hEK8; Tyrosine-protein kinase




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



**Background** 



TYRO1; Tyrosine-protein kinase receptor SEK
This gene belongs to the ephrin receptor subfamily
of the protein-tyrosine kinase family. EPH and
EPH-related receptors have been implicated in
mediating developmental events, particularly in the
nervous system. Receptors in the EPH subfamily
typically have a single kinase domain and an
extracellular region containing a Cys-rich domain
and 2 fibronectin type III repeats. The ephrin
receptors are divided into 2 groups based on the
similarity of their extracellular domain sequences
and their affinities for binding ephrin-A and ephrin-B
ligands. Multiple transcript variants encoding
different isoforms have been found for this gene.
[provided by RefSeq, Jan 2015],

Western Blot analysis of JK cells using Phospho-EphA4 (Y596) Polyclonal Antibody

