



Olfactory receptor 5M3 rabbit pAb

Cat No.:ES5213

For research use only


Overview

Product Name	Olfactory receptor 5M3 rabbit pAb
Host species	Rabbit
Applications	WB;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human OR5M3. AA range:53-102
Specificity	Olfactory receptor 5M3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 5M3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 5M3
Gene Name	OR5M3
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	219482
Human Swiss-Prot Number	Q8NGP4
Alternative Names	OR5M3; Olfactory receptor 5M3; Olfactory receptor OR11-191
Background	olfactory receptor family 5 subfamily M member 3(OR5M3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the







perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Western Blot analysis of HEPG2 cells using Olfactory receptor 5M3 Polyclonal Antibody diluted at 1:1000

Immunofluorescence analysis of MCF7 cells, using OR5M3 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from HepG2, HUVEC, and MCF-7 cells, using OR5M3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR5M3 antibody.

