

DECR2 rabbit pAb

Cat No.:ES5464

For research use only

Overview

Product Name DECR2 rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human DECR2. AA

range:217-266

Specificity DECR2 Polyclonal Antibody detects endogenous

levels of DECR2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Peroxisomal 2,4-dienoyl-CoA reductase

Gene Name DECR2
Cellular localization Peroxisome .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 36kD
Human Gene ID 26063
Human Swiss-Prot Number Q9NUI1

Alternative Names DECR2; PDCR; Peroxisomal 2; 4-dienoyl-CoA

reductase; pDCR; 2,4-dienoyl-CoA reductase 2 catalytic activity:Trans-2,3-didehydroacyl-CoA +

NADP(+) = trans,trans-2,3,4,5-tetradehydroacyl-CoA

+ NADPH., function: Auxiliary enzyme of

beta-oxidation. Participates in the degradation of unsaturated fatty enoyl-CoA esters having double bonds in both even- and odd-numbered positions in




Background

+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com



peroxisome. Catalyzes the NADP-dependent reduction of 2,4-dienoyl-CoA to yield trans-3-enoyl-CoA. Has activity towards short and medium chain 2,4-dienoyl-CoAs, but also towards 2,4,7,10,13,16,19-docosaheptaenoyl-CoA, suggesting that it does not constitute a rate limiting step in the peroxisomal degradation of docosahexaenoic acid.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family. 2,4-dienoyl-CoA reductase subfamily.,



Western blot analysis of lysates from LOVO cells, using DECR2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using DECR2 antibody.

