

## Olfactory receptor 10H3/4 rabbit pAb

Cat No.: ES5500

For research use only

## Overview

Product Name Olfactory receptor 10H3/4 rabbit pAb

Host species Rabbit
Applications WB;ELISA

**Species Cross-Reactivity** Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR10H3/10H4. AA

range:234-283

**Specificity** Olfactory receptor 10H3/4 Polyclonal Antibody

detects endogenous levels of Olfactory receptor

10H3/4 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 10H3/4

Gene Name OR10H3/OR10H4

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band38kD

Human Gene ID 26532/126541 Human Swiss-Prot Number 060404/Q8NGA5

Alternative Names OR10H3; Olfactory receptor 10H3; Olfactory

receptor OR19-24; OR10H4; Olfactory receptor

10H4; Olfactory receptor OR19-28

**Background** Olfactory receptors interact with odorant molecules


in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of




+86-27-59760950 ELKbio@ELKbiotech.com www.elk



G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western Blot analysis of HELA using Olfactory receptor 10H3/4 Polyclonal Antibody. Antibody was diluted at 1:1000



Western blot analysis of lysates from NIH/3T3 cells, using OR10H3/10H4 Antibody. The lane on the right is blocked with the synthesized peptide.

