

Olfactory receptor 8B2/3 rabbit pAb

Cat No.:ES5503

For research use only

Overview

Product Name	Olfactory receptor 8B2/3 rabbit pAb
Host species	Rabbit
Applications	WB;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunofluorescence:
	1/200 - 1/1000. ELISA: 1/10000. Not yet tested in
	other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human OR8B2/8B3. AA
	range:199-248
Specificity	Olfactory receptor 8B2/3 Polyclonal Antibody
	detects endogenous levels of Olfactory receptor
	8B2/3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 8B2/3
Gene Name	OR8B2/OR8B3
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	26595
Human Swiss-Prot Number	Q96RD0/Q8NGG8
Alternative Names	OR8B2; Olfactory receptor 8B2; Olfactory receptor
	OR11-309; OR8B3; Olfactory receptor 8B3; Olfactory
	receptor OR11-311
Background	olfactory receptor family 8 subfamily B member
	2(OR8B2) Homo sapiens Olfactory receptors
	interact with odorant molecules in the nose, to



+86-27-59760950


ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from Jurkat and HT-29 -- 117 cells, using OR8B2/8B3 Antibody. The lane on the right is -- 85 blocked with the synthesized peptide.

Western blot analysis of the lysates from K562 cells using OR8B2/B3 antibody.



ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C