

Dynein IC1 rabbit pAb

Cat No.:ES5529

For research use only

Overview

| Product Name | Dynein IC1 rabbit pAb |
|--------------------------|--|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Mouse;Rat |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not |
| | yet tested in other applications. |
| Immunogen | The antiserum was produced against synthesized |
| | peptide derived from human DNAI1. AA |
| | range:211-260 |
| Specificity | Dynein IC1 Polyclonal Antibody detects endogenous |
| | levels of Dynein IC1 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Dynein intermediate chain 1 axonemal |
| Gene Name | DNAI1 |
| Cellular localization | Dynein axonemal particle . Cytoplasm, cytoskeleton, |
| | cilium axoneme . |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 79kD |
| Human Gene ID | 27019 |
| Human Swiss-Prot Number | Q9UI46 |
| Alternative Names | DNAI1; Dynein intermediate chain 1; axonemal; |
| | Axonemal dynein intermediate chain 1 |
| Background | This gene encodes a member of the dynein |
| | intermediate chain family. The encoded protein is |
| | part of the dynein complex in respiratory cilia. The |
| | inner- and outer-arm dyneins, which bridge |
| | between the doublet microtubules in axonemes, are |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



the force-generating proteins responsible for the sliding movement in axonemes. The intermediate and light chains, thought to form the base of the dynein arm, help mediate attachment and may also participate in regulating dynein activity. Mutations in this gene result in abnormal ciliary ultrastructure and function associated with primary ciliary dyskinesia and Kartagener syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],

Western blot analysis of lysates from COLO cells, using DNAI1 Antibody. The lane on the right is blocked with the synthesized peptide.





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C