

Olfactory receptor 5W2 rabbit pAb

Cat No.:ES6055

For research use only

Overview

| Product Name | Olfactory receptor 5W2 rabbit pAb |
|------------------------------|---|
| Host species | Rabbit |
| Applications | WB;IF;ELISA |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. Immunofluorescence: |
| | 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in |
| | other applications. |
| Immunogen | The antiserum was produced against synthesized |
| _ | peptide derived from human OR5W2. AA |
| | range:158-207 |
| Specificity | Olfactory receptor 5W2 Polyclonal Antibody detects |
| | endogenous levels of Olfactory receptor 5W2 |
| | protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Olfactory receptor 5W2 |
| Gene Name | OR5W2 |
| Cellular localization | Cell membrane; Multi-pass membrane protein. |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 34kD |
| Human Gene ID | 390148 |
| Human Swiss-Prot Number | Q8NH69 |
| Alternative Names | OR5W2; OR5W2P; OR5W3P; Olfactory receptor |
| | 5W2; Olfactory receptor 5W3; Olfactory receptor |
| | OR11-155 |
| Background | olfactory receptor family 5 subfamily W member |
| | 2(OR5W2) Homo sapiens Olfactory receptors |
| | interact with odorant molecules in the nose, to |
| | |



+86-27-59760950


ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],





Immunofluorescence analysis of A549 cells, using OR5W2 Antibody. The picture on the right is blocked with the synthesized peptide.






+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from HeLa cells, using OR5W2 Antibody. The lane on the right is blocked with the synthesized peptide.

Western blot analysis of the lysates from 293 cells using OR5W2 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C