

Olfactory receptor 10K1/2 rabbit pAb

Cat No.: ES6078

For research use only

Overview

Product Name Olfactory receptor 10K1/2 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/20000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR10K1/10K2. AA

range:56-105

Specificity Olfactory receptor 10K1/2 Polyclonal Antibody

detects endogenous levels of Olfactory receptor

10K1/2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 10K1/2

Gene Name OR10K1/OR10K2

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band35kD

Human Gene ID 391109/391107 **Human Swiss-Prot Number** Q8NGX5/Q6IF99

Alternative Names OR10K1; Olfactory receptor 10K1; Olfactory

receptor OR1-6; OR10K2; Olfactory receptor 10K2;

Olfactory receptor OR1-4

Background Olfactory receptors interact with odorant molecules


in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory






receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Immunofluorescence analysis of LOVO cells, using OR10K1/10K2 Antibody. The picture on the right is blocked with the synthesized peptide.




-- 117
Western blot analysis of lysates from HepG2 cells, using
OR10K1/10K2 Antibody. The lane on the right is blocked
with the synthesized peptide.



+86-27-59760950







Western blot analysis of the lysates from HUVECcells using OR10K1/10K2 antibody.

