

Tau (phospho Ser519) rabbit pAb

Cat No.:ES6178

For research use only

Overview

| Product Name | Tau (phospho Ser519) rabbit pAb |
|--------------------------|---|
| Host species | Rabbit |
| Applications | WB;ELISA |
| Species Cross-Reactivity | Human;Mouse;Rat |
| Recommended dilutions | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not |
| | yet tested in other applications. |
| Immunogen | The antiserum was produced against synthesized |
| - | peptide derived from human Tau around the |
| | phosphorylation site of Ser519/202. AA |
| | range:266-315 |
| Specificity | Phospho-Tau (S519) Polyclonal Antibody detects |
| | endogenous levels of Tau protein only when |
| | phosphorylated at S519. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Microtubule-associated protein tau |
| Gene Name | MAPT |
| Cellular localization | Cytoplasm, cytosol . Cell membrane ; Peripheral |
| | membrane protein ; Cytoplasmic side . Cytoplasm, |
| | cytoskeleton . Cell projection, axon . Cell projection, |
| | dendrite . Secreted . Mostly found in the axons of |
| | neurons, in the cytosol and in association with pla |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 50-85kD |
| Human Gene ID | 4137 |
| Human Swiss-Prot Number | P10636 |
| Alternative Names | MAPT; MAPTL; MTBT1; TAU; Microtubule-associated |
| | protein tau; Neurofibrillary tangle protein; Paired |



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

helical filament-tau; PHF-tau

This gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from Jurkat cells treated with H2O2 100uM 30', using Tau (Phospho-Ser519/202) Antibody. The lane on the right is blocked with the phospho peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C