

## **Olfactory receptor 13G1 rabbit pAb**

## Cat No.:ES6275

For research use only

## Overview

Product Name	Olfactory receptor 13G1 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human OR13G1. AA
	range:241-290
Specificity	Olfactory receptor 13G1 Polyclonal Antibody detects
	endogenous levels of Olfactory receptor 13G1
	protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 13G1
Gene Name	OR13G1
<b>Cellular localization</b>	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	441933
Human Swiss-Prot Number	Q8NGZ3
Alternative Names	OR13G1; Olfactory receptor 13G1; Olfactory
	receptor OR1-37
Background	Olfactory receptors interact with odorant molecules
	in the nose, to initiate a neuronal response that
	triggers the perception of a smell. The olfactory
	receptor proteins are members of a large family of
	G-protein-coupled receptors (GPCR) arising from



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of 293 cells using Olfactory receptor 13G1 Polyclonal Antibody




Western blot analysis of lysates from 293 cells, using OR13G1 Antibody. The lane on the right is blocked with the synthesized peptide.



www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of the lysates from Jurkat cells using OR13G1 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C