

## NTR1 rabbit pAb

Cat No.:ES6435

For research use only

## Overview

**Product Name** NTR1 rabbit pAb

**Host species** Rabbit

WB;IHC;IF;ELISA **Applications Species Cross-Reactivity** Human; Mouse

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

**Immunogen** The antiserum was produced against synthesized

peptide derived from human NTR1. AA

range:181-230

**Specificity** NTR1 Polyclonal Antibody detects endogenous

levels of NTR1 protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. Storage

**Protein Name** Neurotensin receptor type 1

**Gene Name** NTSR1

Cellular localization Cell membrane; Multi-pass membrane protein.

> Membrane raft. Palmitoylation is required for localization at CAV1-enriched membrane rafts. . The antibody was affinity-purified from rabbit

Purification

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 46kD **Human Gene ID** 4923 **Human Swiss-Prot Number** P30989

**Alternative Names** NTSR1; NTRR; Neurotensin receptor type 1; NT-R-1;

NTR1; High-affinity levocabastine-insensitive

neurotensin receptor; NTRH


Background Neurotensin receptor 1 belongs to the large



+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com





superfamily of G-protein coupled receptors. NTSR1 mediates the multiple functions of neurotensin, such as hypotension, hyperglycemia, hypothermia, antinociception, and regulation of intestinal motility and secretion. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from MCF7 cells, using NTR1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

