

ATP5G3 rabbit pAb

Cat No.: ES6568

For research use only

Overview

Product Name ATP5G3 rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA
Species Cross-Reactivity Human;Rat

Recommended dilutions Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human ATP5G3. AA range:1-50

Specificity ATP5G3 Polyclonal Antibody detects endogenous

levels of ATP5G3 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name ATP synthase lipid-binding protein mitochondrial

Gene Name ATP5G3

Cellular localization Mitochondrion membrane; Multi-pass membrane

protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Human Gene ID 518 Human Swiss-Prot Number P48201

Alternative Names ATP5G3; ATP synthase lipid-binding protein;

mitochondrial; ATP synthase proteolipid P3; ATPase

protein 9; ATPase subunit c

Background This gene encodes a subunit of mitochondrial ATP

synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech




phosphorylation. ATP synthase is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene is one of three genes that encode subunit c of the proton channel. Each of the three genes have distinct mitochondrial import sequences but encode the identi

Immunofluorescence analysis of A549 cells, using ATP5G3 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human pancreas tissue, using ATP5G3 Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950

