

PR (phospho Ser294) rabbit pAb

Cat No.: ES6579

For research use only

Overview

Product Name PR (phospho Ser294) rabbit pAb

Host species Rabbit
Applications WB;IHC

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300

Immunogen The antiserum was produced against synthesized

peptide derived from human Progesterone Receptor

around the phosphorylation site of Ser294. AA

range:261-310

Specificity Phospho-PR (S294) Polyclonal Antibody detects

endogenous levels of PR protein only when

phosphorylated at S294.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Progesterone receptor

Gene Name PGR

Cellular localization Nucleus. Cytoplasm. Nucleoplasmic shuttling is both

hormone- and cell cycle-dependent. On hormone stimulation, retained in the cytoplasm in the G(1)

and G(2)/M phases.; [Isoform A]: Nucleus. Cytoplasm. Mainly nuclear.; [Isoform 4]:

Mitochondrion outer membrane.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band98kDHuman Gene ID5241Human Swiss-Prot NumberP06401

Alternative Names PGR; NR3C3; Progesterone receptor; PR; Nuclear

receptor subfamily 3 group C member 3




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




Background

This gene encodes a member of the steroid receptor superfamily. The encoded protein mediates the physiological effects of progesterone, which plays a central role in reproductive events associated with the establishment and maintenance of pregnancy. This gene uses two distinct promotors and translation start sites in the first exon to produce several transcript variants, both protein coding and non-protein coding. Two of the isoforms (A and B) are identical except for an additional 165 amino acids found in the N-terminus of isoform B and mediate their own response genes and physiologic effects with little overlap. [provided by RefSeq, Sep 2015],



Western Blot analysis of 293 cells using Phospho-PR (S294) Polyclonal Antibody





Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Progesterone Receptor (Phospho-Ser294) Antibody





Immunofluorescence analysis of HeLa cells, using Progesterone Receptor (Phospho-Ser294) Antibody. The picture on the right is blocked with the phospho peptide.





Western blot analysis of lysates from 293 cells treated with Etoposide 25uM 60', using Progesterone Receptor (Phospho-Ser294) Antibody. The lane on the right is blocked with the phospho peptide.

