



FGFR-5 rabbit pAb

Cat No.:ES6628

For research use only


Overview

Product Name	FGFR-5 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	Synthesized peptide derived from FGFR-5 . at AA range: 130-210
Specificity	FGFR-5 Polyclonal Antibody detects endogenous levels of FGFR-5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Fibroblast growth factor receptor-like 1
Gene Name	FGFRL1
Cellular localization	Membrane ; Single-pass type I membrane protein . Predominantly localized in the plasma membrane but also detected in the Golgi and in secretory vesicles.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	54kD
Human Gene ID	53834
Human Swiss-Prot Number	Q8N441
Alternative Names	FGFRL1; FGFR5; FHFR; Fibroblast growth factor receptor-like 1; FGF receptor-like protein 1; FGF homologous factor receptor; FGFR-like protein; Fibroblast growth factor receptor 5; FGFR-5
Background	The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family,





where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. A marked difference between this gene product and the other family members is its lack of a cytoplasmic tyrosine kinase domain. The result is a transmembrane receptor that could interact with other



Western Blot analysis of 3T3 cells using FGFR-5 Polyclonal Antibody

