

## IPP-2 (phospho Ser44) rabbit pAb

## Cat No.:ES6695

For research use only

## Overview

Product Name	IPP-2 (phospho Ser44) rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human; Mouse; Rat
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/40000. Not yet tested in other applications.
Immunogen	Synthesized phospho-peptide around the
	phosphorylation site of human IPP-2 (phospho
	Ser44)
Specificity	Phospho-IPP-2 (S44) Polyclonal Antibody detects
	endogenous levels of IPP-2 protein only when
	phosphorylated at S44.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Protein phosphatase inhibitor 2
Gene Name	PPP1R2
<b>Cellular localization</b>	protein phosphatase type 1 complex,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	44kD
Human Gene ID	5504
Human Swiss-Prot Number	P41236
Alternative Names	PPP1R2; IPP2; Protein phosphatase inhibitor 2; IPP-2
Background	Protein phosphatase-1 (PP1) is one of the main
	eukaryotic serine/threonine phosphatases. The
	protein encoded by this gene binds to the catalytic
	subunit of PP1, strongly inhibiting its activity. Ten
	related pseudogenes have been found throughout



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



the human genome. Several splice variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],

Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C