

PKD1/2/3 (phospho Ser738/S742) rabbit pAb

Cat No.: ES6789

For research use only

Overview

Product Name PKD1/2/3 (phospho Ser738/S742) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human PKD1/2/3/PKC mu around the phosphorylation site of Ser738 and

Ser742. AA range:706-755

Specificity Phospho-PKD1/2/3 (S738/S742) Polyclonal Antibody

detects endogenous levels of PKD1/2/3 protein only

when phosphorylated at S738/S742.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/threonine-protein kinase D1/2/3

Gene Name KPCD1/KPCD2/KPCD3

Cellular localization Cytoplasm . Cell membrane . Golgi apparatus,

trans-Golgi network. Translocation to the cell membrane is required for kinase activation.

The antihody was affinity-purified from rabbit

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band101kD

Human Gene ID 5587/25865/23683
Human Swiss-Prot Number Q15139/Q9BZL6/O94806
Alternative Names PRKD1; PKD; PKD1; PRKCM;

Serine/threonine-protein kinase D1; Protein kinase




+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com




Background

C mu type; Protein kinase D; nPKC-D1; nPKC-mu; PRKD2; PKD2; HSPC187; Serine/threonine-protein kinase D2; nPKC-D2; PRKD3; EPK2; PRKCN; Serine/threonine-protein kinas PRKD1 is a serine/threonine kinase that regulates a variety of cellular functions, including membrane receptor signaling, transport at the Golgi, protection from oxidative stress at the mitochondria, gene transcription, and regulation of cell shape, motility, and adhesion (summary by Eiseler et al., 2009 [PubMed 19329994]).[supplied by OMIM, Nov 2010],



Western Blot analysis of A549 cells using Phospho-PKD1/2/3 (S738/S742) Polyclonal Antibody




+86-27-59760950


Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PKD1/2/3/PKC mu (Phospho-Ser738+Ser742) Antibody







Immunohistochemistry analysis of paraffin-embedded human heart, using PKD1/2/3/PKC mu (Phospho-Ser738+Ser742) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from A549 cells treated with PMA 125ng/ml 30' and HT29 cells treated with serum 20% 15', using PKD1/2/3/PKC mu (Phospho-Ser738+Ser742) Antibody. The lane on the right is blocked with the phospho peptide.

