

## Rb (phospho Ser788) rabbit pAb

Cat No.: ES6998

For research use only

## Overview

Product Name Rb (phospho Ser788) rabbit pAb

Host species Rabbit

Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

**Recommended dilutions** Immunohistochemistry: 1/100 - 1/300. ELISA:

1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human Retinoblastoma around the phosphorylation site of Ser788. AA

range:756-805

Specificity Phospho-Rb (S788) Polyclonal Antibody detects

endogenous levels of Rb protein only when

phosphorylated at S788.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Retinoblastoma-associated protein

Gene Name RB1

Cellular localization Nucleus . During keratinocyte differentiation,

acetylation by KAT2B/PCAF is required for nuclear

localization. .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 5925 Human Swiss-Prot Number P06400

Alternative Names RB1; Retinoblastoma-associated protein; p105-Rb;

pRb; Rb; pp110

**Background** The protein encoded by this gene is a negative

regulator of the cell cycle and was the first tumor



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiot



suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma. [provided by RefSeq, Jul 2008],



+86-27-59760950

Immunohistochemical analysis of paraffin-embedded human Small intestinal stromal tumor. 1,
Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).

