



NDRG4 rabbit pAb

Cat No.:ES7211

For research use only

Overview

Product Name	NDRG4 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human NDRG4. AA range:187-236
Specificity	NDRG4 Polyclonal Antibody detects endogenous levels of NDRG4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Protein NDRG4
Gene Name	NDRG4
Cellular localization	Cytoplasm, cytosol .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	34kD
Human Gene ID	65009
Human Swiss-Prot Number	Q9ULP0
Alternative Names	NDRG4; BDM1; KIAA1180; Protein NDRG4; Brain development-related molecule 1; N-myc downstream-regulated gene 4 protein; Vascular smooth muscle cell-associated protein 8; SMAP-8
Background	This gene is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. The protein encoded by this








gene is a cytoplasmic protein that is required for cell cycle progression and survival in primary astrocytes and may be involved in the regulation of mitogenic signalling in vascular smooth muscles cells.

Alternative splicing results in multiple transcripts encoding different isoforms.[provided by RefSeq, Jun 2011],

Immunohistochemistry analysis of paraffin-embedded human heart tissue, using NDRG4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HT-29 and HepG2 cells, using NDRG4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using NDRG4 antibody.

