

Raf-B (phospho Ser602) rabbit pAb

Cat No.: ES7279

For research use only

Overview

Product Name Raf-B (phospho Ser602) rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human B-RAF around the phosphorylation site of Ser602. AA range:576-625

Specificity Phospho-Raf-B (S602) Polyclonal Antibody detects

endogenous levels of Raf-B protein only when

phosphorylated at S602.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Serine/threonine-protein kinase B-raf

Gene Name BRAF

Cellular localization Nucleus . Cytoplasm . Cell membrane . Colocalizes

with RGS14 and RAF1 in both the cytoplasm and

membranes...

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 85kD
Human Gene ID 673
Human Swiss-Prot Number P15056

Alternative Names BRAF; BRAF1; RAFB1; Serine/threonine-protein

kinase B-raf; Proto-oncogene B-Raf; p94; v-Raf murine sarcoma viral oncogene homolog B1


Background This gene encodes a protein belonging to the raf/mil




+86-27-59760950 ELKbio@ELKbiotech.com



family of serine/threonine protein kinases. This protein plays a role in regulating the MAP kinase/ERKs signaling pathway, which affects cell division, differentiation, and secretion. Mutations in this gene are associated with cardiofaciocutaneous syndrome, a disease characterized by heart defects, mental retardation and a distinctive facial appearance. Mutations in this gene have also been associated with various cancers, including non-Hodgkin lymphoma, colorectal cancer, malignant melanoma, thyroid carcinoma, non-small cell lung carcinoma, and adenocarcinoma of lung. A pseudogene, which is located on chromosome X, has been identified for this gene. [provided by RefSeq, Jul 2008],



Western Blot analysis of 3T3 cells using Phospho-Raf-B (S602) Polyclonal Antibody




+86-27-59760950

Immunohistochemistry analysis of paraffin-embedded human lymph node, using B-RAF (Phospho-Ser602) Antibody. The picture on the right is blocked with the phospho peptide.







Western blot analysis of lysates from NIH/3T3 cells treated with EGF 200ng/ml 30', using B-RAF (Phospho-Ser602) Antibody. The lane on the right is blocked with the phospho peptide.



+86-27-59760950