

## Olfactory receptor 6J1 rabbit pAb

Cat No.: ES7579

For research use only

## Overview

Product Name Olfactory receptor 6J1 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR6J1. AA

range:231-280

**Specificity** Olfactory receptor 6J1 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 6J1 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 6J1

Gene Name OR6J1

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 39kD
Human Gene ID 79549
Human Swiss-Prot Number Q8NGC5

Alternative Names OR6J1; OR6J2; Olfactory receptor 6J1; Olfactory

receptor 6J2

**Background** Olfactory receptors interact with odorant molecules

in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from





+86-27-59760950 ELKbio@ELKbiotech.com v




single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Immunofluorescence analysis of MCF7 cells, using OR6J1 Antibody. The picture on the right is blocked with the synthesized peptide.








Western blot analysis of lysates from HUVEC and MCF-7 cells, using OR6J1 Antibody. The lane on the right is blocked with the synthesized peptide.







Western blot analysis of the lysates from COLO205 cells using OR6J1 antibody.

