

FGF-23 rabbit pAb

Cat No.: ES7620

For research use only

Overview

Product Name FGF-23 rabbit pAb

Host species Rabbit

Applications WB;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300

ELISA: 1/20000. IF 1:100-300 Not yet tested in other

applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human FGF23. AA

range:151-200

Specificity FGF-23 Polyclonal Antibody detects endogenous

levels of FGF-23 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Fibroblast growth factor 23

Gene Name FGF23

Cellular localization Secreted . Secretion is dependent on

O-glycosylation.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 27kD
Human Gene ID 8074
Human Swiss-Prot Number Q9GZV9

Alternative Names FGF23; HYPF; Fibroblast growth factor 23; FGF-23;

Phosphatonin; Tumor-derived hypophosphatemia-inducing factor


Background This gene encodes a member of the fibroblast

growth factor family of proteins, which possess broad mitogenic and cell survival activities and are







involved in a variety of biological processes. The product of this gene regulates phosphate homeostasis and transport in the kidney. The full-length, functional protein may be deactivated via cleavage into N-terminal and C-terminal chains. Mutation of this cleavage site causes autosomal dominant hypophosphatemic rickets (ADHR). Mutations in this gene are also associated with hyperphosphatemic familial tumoral calcinosis (HFTC). [provided by RefSeq, Feb 2013],



Immunofluorescence analysis of A549. 1, primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.




Immunofluorescence analysis of HUVEC cells, using FGF23 Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950







Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night

