

Olfactory receptor 51G2 rabbit pAb

Cat No.: ES7632

For research use only

Overview

Product Name Olfactory receptor 51G2 rabbit pAb

Host species Rabbit WB;ELISA **Applications**

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

The antiserum was produced against synthesized **Immunogen**

peptide derived from human OR51G2. AA

range:211-260

Specificity Olfactory receptor 51G2 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 51G2

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles. Storage

Olfactory receptor 51G2 **Protein Name**

Gene Name OR51G2

Cellular localization Cell membrane; Multi-pass membrane protein. Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 35kD **Human Gene ID** 81282 **Human Swiss-Prot Number** Q8NGK0

Alternative Names OR51G2; Olfactory receptor 51G2; Olfactory

receptor OR11-28

Olfactory receptors interact with odorant molecules **Background**


> in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from




+86-27-59760950 ELKbio@ELKbiotech.com



single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



Western blot analysis of lysates from HeLa and HepG2 cells, using OR51G2 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

Western blot analysis of the lysates from COLO205 cells using OR51G2 antibody.

