

Caspase-2 (phospho Ser157) rabbit pAb

Cat No.: ES7685

For research use only

Overview

Product Name Caspase-2 (phospho Ser157) rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Immunohistochemistry: 1/100 - 1/300. ELISA:

1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human Caspase 2 around the phosphorylation site of Ser157. AA range:123-172

Specificity Phospho-Caspase-2 (S157) Polyclonal Antibody

detects endogenous levels of Caspase-2 protein only

when phosphorylated at S157.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Caspase2
Gene Name CASP2

Cellular localization nucleus, cytoplasm, mitochondrion, cytosol, membrane,

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Background

Human Gene ID 835 Human Swiss-Prot Number P42575

Alternative Names CASP2; ICH1; NEDD2; Caspase-2; CASP-2; Neural

precursor cell expressed developmentally

down-regulated protein 2; NEDD-2; Protease ICH-1
This gene encodes a member of the cysteine-aspartic

acid protease (caspase) family. Caspases mediate cellular apoptosis through the proteolytic cleavage of


specific protein substrates. The encoded protein may






function in stress-induced cell death pathways, cell cycle maintenance, and the suppression of tumorigenesis. Increased expression of this gene may play a role in neurodegenerative disorders including Alzheimer's disease, Huntington's disease and temporal lobe epilepsy. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2011],

Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbe



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using Caspase 2 (Phospho-Ser157) Antibody. The picture on the right is blocked with the phospho peptide.



+86-27-59760950

