

MAPKAPK-2 rabbit pAb

Cat No.:ES8007

For research use only

Overview

Product Name	MAPKAPK-2 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/5000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human MAPKAPK2. AA
	range:238-287
Specificity	MAPKAPK-2 Polyclonal Antibody detects
	endogenous levels of MAPKAPK-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	MAP kinase-activated protein kinase 2
Gene Name	МАРКАРК2
Cellular localization	Cytoplasm . Nucleus . Phosphorylation and
	subsequent activation releases the autoinhibitory
	helix, resulting in the export from the nucleus into
	the cytoplasm.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	46kD
Human Gene ID	9261
Human Swiss-Prot Number	P49137
Alternative Names	MAPKAPK2; MAP kinase-activated protein kinase 2;
	MAPK-activated protein kinase 2; MAPKAP kinase 2;
	МАРКАР-К2; МАРКАРК-2; МК-2; МК2
Background	This gene encodes a member of the Ser/Thr protein



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



kinase family. This kinase is regulated through direct phosphorylation by p38 MAP kinase. In conjunction with p38 MAP kinase, this kinase is known to be involved in many cellular processes including stress and inflammatory responses, nuclear export, gene expression regulation and cell proliferation. Heat shock protein HSP27 was shown to be one of the substrates of this kinase in vivo. Two transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MAPKAPK2 Antibody. The picture on the right is blocked with the synthesized peptide.






Western blot analysis of lysates from COS cells, using MAPKAPK2 Antibody. The lane on the right is blocked with the synthesized peptide.



ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of the lysates from HeLa cells using MAPKAPK2 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C