

Cdc25A (phospho Ser75) rabbit pAb

Cat No.:ES8127

For research use only

Overview

Product Name Cdc25A (phospho Ser75) rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not

yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human CDC25A around the phosphorylation site of Ser75. AA range:43-92 Phospho-Cdc25A (S75) Polyclonal Antibody detects

Specificity Phospho-Cdc25A (\$75) Polyclonal Antibody detects

endogenous levels of Cdc25A protein only when

phosphorylated at S75.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name M-phase inducer phosphatase 1

Gene Name CDC25A

Cellular localization intracellular, nucleus, nucleoplasm, cytoplasm, cytosol,

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 60kD
Human Gene ID 993
Human Swiss-Prot Number P30304

Alternative Names CDC25A; M-phase inducer phosphatase 1; Dual

specificity phosphatase Cdc25A

Background cell division cycle 25A(CDC25A) Homo sapiens

CDC25A is a member of the CDC25 family of


phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two




+86-27-59760950 ELKbio@ELKbiotech.com www.elkl



phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an oncogene, although its exact role in oncogenesis has not been demonstrated. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],





CDC25A----(pSer75) -- 48
-- 34
-- 26
-- 19
(kD)

Western blot analysis of lysates from Jurkat cells treated with UV, using CDC25A (Phospho-Ser75) Antibody. The lane on the right is blocked with the phospho peptide.

