



# ADAMTS-2 rabbit pAb

Cat No.:ES8355

For research use only


## Overview

<b>Product Name</b>	ADAMTS-2 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
<b>Immunogen</b>	Synthesized peptide derived from ADAMTS-2 . at AA range: 1140-1220
<b>Specificity</b>	ADAMTS-2 Polyclonal Antibody detects endogenous levels of ADAMTS-2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	A disintegrin and metalloproteinase with thrombospondin motifs 2
<b>Gene Name</b>	ADAMTS2
<b>Cellular localization</b>	Secreted, extracellular space, extracellular matrix .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	100kD
<b>Human Gene ID</b>	9509
<b>Human Swiss-Prot Number</b>	O95450
<b>Alternative Names</b>	ADAMTS2; PCINP; PCPNI; A disintegrin and metalloproteinase with thrombospondin motifs 2; ADAM-TS 2; ADAM-TS2; ADAMTS-2; Procollagen I N-proteinase; PC I-NP; Procollagen I/II amino propeptide-processing enzyme; Procollagen N-endopeptidase; pNPI
<b>Background</b>	This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with





thrombospondin motifs) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The encoded preproprotein is proteolytically processed to generate the mature procollagen N-proteinase. This proteinase excises the N-propeptide of the fibrillar procollagens types I-III and type V. Mutations in this gene cause Ehlers-Danlos syndrome type VIIC, a recessively inherited connective-tissue disorder. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically



Western blot analysis of HepG2 K562 HeLa Colo using ADAMTS-2 antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

