

## EDA rabbit pAb

## Cat No.:ES8383

For research use only

## Overview

Product Name	EDA rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300.
	ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from the Internal region of human
	EDA. AA range:120-170
Specificity	EDA Polyclonal Antibody detects endogenous levels
	of EDA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Ectodysplasin-A
Gene Name	EDA
Cellular localization	Cell membrane ; Single-pass type II membrane
	protein .; [Ectodysplasin-A, secreted form]:
	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	42kD
Human Gene ID	1896
Human Swiss-Prot Number	Q92838
Alternative Names	EDA; ED1; EDA2; Ectodysplasin-A; Ectodermal
	dysplasia protein; EDA protein
Background	The protein encoded by this gene is a type II
	membrane protein that can be cleaved by furin to
	produce a secreted form. The encoded protein,
	which belongs to the tumor necrosis factor family,



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



138 100

70-55-


40-35--25--

15--

acts as a homotrimer and may be involved in cell-cell signaling during the development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia, anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Western blot analysis of mouse-lung mouse-heart mouse-liver lysis using EDA antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

Immunohistochemical analysis of paraffin-embedded human-breast-cancer, antibody was diluted at 1:200



EDA



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C