

CD249 rabbit pAb

Cat No.: ES8413

For research use only

Overview

Product Name CD249 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA **Species Cross-Reactivity** Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300.

ELISA: 1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from the Internal region of human

ENPEP. AA range:671-720

Specificity CD249 Polyclonal Antibody detects endogenous

levels of CD249 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Glutamyl aminopeptidase

Gene Name ENPEP

Cell ular localization Cell membrane ; Single-pass type II membrane

protein.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 110kD
Human Gene ID 2028
Human Swiss-Prot Number Q07075

Alternative Names ENPEP; Glutamyl aminopeptidase; EAP;

Aminopeptidase A; AP-A; Differentiation antigen

gp160; CD249

Background catalytic activity:Release of N-terminal glutamate

(and to a lesser extent aspartate) from a peptide.,cofactor:Binds 1 zinc ion per

subunit., function: Appears to have a role in the







catabolic pathway of the renin-angiotensin system. Probably plays a role in regulating growth and differentiation of early B-lineage cells.,similarity:Belongs to the peptidase M1 family.,subunit:Homodimer; disulfide-linked.,tissue specificity:Expressed by epithelial cells of the proximal tubule cells and the glomerulus of the nephron. Also found in a variety of other tissues.,

MCF-7
178-100-70-55-40-35-25--

Western Blot analysis of MCF7 cells using CD249 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded rat-kidney, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded rat-kidney, antibody was diluted at 1:200



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com