

HNF-4α (Acetyl Lys106) rabbit pAb

Cat No.: ES8628

For research use only

Overview

Product Name HNF-4α (Acetyl Lys106) rabbit pAb

Host species Rabbit
Applications WB;ELISA

Species Cross-Reactivity Human; Rat; Mouse

Recommended dilutions WB 1:500-2000, ELISA 1:10000-20000

Immunogen Synthetic Acetyl peptide from human protein at AA

range: 106

Specificity The antibody detects endogenous HNF- 4α when

Acetyl occurs at Lys106

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Hepatocyte nuclear factor 4-alpha (HNF-4-alpha)

(Nuclear recentor subfamily 2 group A member 1)

(Nuclear receptor subfamily 2 group A member 1) (Transcription factor 14) (TCF-14) (Transcription

factor HNF-4)

Gene Name HNF4A HNF4 NR2A1 TCF14

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 55kD
Human Gene ID 3172
Human Swiss-Prot Number P41235

Alternative Names Hepatocyte nuclear factor 4-alpha (HNF-4-alpha)

(Nuclear receptor subfamily 2 group A member 1) (Transcription factor 14) (TCF-14) (Transcription

factor HNF-4)


Background The protein encoded by this gene is a nuclear

transcription factor which binds DNA as a homodimer. The encoded protein controls the



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.co





+86-27-59760950

expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This gene may play a role in development of the liver, kidney, and intestines. Mutations in this gene have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I. Alternative splicing of this gene results in multiple transcript variants encoding several different isoforms. [provided by RefSeq, Apr 2012],

Western blot analysis of mouse-lung mouse-kidney mouse-liver lysate, antibody was diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

