

Annexin I rabbit pAb

Cat No.:ES8636

For research use only

Overview

Product Name	Annexin I rabbit pAb	
Host species	Rabbit	
Applications	IF;WB;IHC;ELISA	
Species Cross-Reactivity	Human;Rat;Mouse;	
Recommended dilutions	IF: 1:50-200 WB 1:500-2000,IHC-p 1:500-200, ELISA	
	1:10000-20000	
Immunogen	Synthetic peptide from human protein at AA range:	
	130-180	
Specificity	The antibody detects endogenous Annexin I	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	Annexin A1 (Annexin I) (Annexin-1) (Calpactin II)	
	(Calpactin-2) (Chromobindin-9) (Lipocortin I)	
	(Phospholipase A2 inhibitory protein) (p35)	
Gene Name	ANXA1 ANX1 LPC1	
Cellular localization	Nucleus . Cytoplasm . Cell projection, cilium . Cell	
	membrane . Membrane ; Peripheral membrane	
	protein . Endosome membrane ; Peripheral	
	membrane protein . Basolateral cell membrane .	
	Apical cell membrane . Lateral cell membrane .	
	Secreted . Secreted, extracellular space . Cell	
	membrane ; Peripheral membrane protein ;	
	Extracellular side . Secreted, extracellular exosome .	
	Cytoplasmic vesicle, secretory vesicle lumen . Cell	
	projection, phagocytic cup . Early endosome .	
	Cytoplasmic vesicle membrane ; Peripheral	
	membrane protein . Secreted, at least in part via	
	exosomes and other secretory vesicles. Detected in	
	exosomes and other extracellular vesicles	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
	(PubMed:25664854). Alternatively, the secretion is	
	dependent on protein unfolding and facilitated by	
	the cargo receptor TMED10; it results in t	



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C




Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	38kD
Human Gene ID	301
Human Swiss-Prot Number	P04083
Alternative Names	Annexin A1 (Annexin I) (Annexin-1) (Calpactin II)
	(Calpactin-2) (Chromobindin-9) (Lipocortin I)
	(Phospholipase A2 inhibitory protein) (p35)
Background	This gene encodes a membrane-localized protein
	that binds phospholipids. This protein inhibits
	phospholipase A2 and has anti-inflammatory
	activity. Loss of function or expression of this gene
	has been detected in multiple tumors. [provided by
	RefSeq, Dec 2014],

Immunofluorescence analysis of human-breast tissue. 1,Annexin I Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target.



Immunofluorescence analysis of human-breast tissue. 1,Annexin I Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Immunohistochemical analysis of paraffin-embedded Human-breast-cancer tissue. 1,Annexin I Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1



Immunohistochemical analysis of paraffin-embedded Human-liver-cancer tissue. 1,Annexin I Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room tempeRature, 30min). Negative control was used by secondary antibody only.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C