

pS2 rabbit pAb

Cat No.:ES8664

For research use only

Overview

Product Name pS2 rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions IHC-p 1:50-200, ELISA 1:10000-20000

Immunogen Synthetic peptide from human protein at AA range:

10-50

Specificity The antibody detects endogenous pS2

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Trefoil factor 1 (Breast cancer estrogen-inducible

protein) (PNR-2) (Polypeptide P1.A) (hP1.A) (Protein

pS2)

Gene Name TFF1 BCEI PS2
Cellular localization Secreted .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band

Human Gene ID 7031 Human Swiss-Prot Number P04155

Alternative Names Trefoil factor 1 (Breast cancer estrogen-inducible

protein;PNR-2;Polypeptide P1.A;hP1.A;Protein pS2) Members of the trefoil family are characterized by

Background Members of the trefoil family are characterized by

having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfides. They are stable secretory proteins expressed in gastrointestinal mucosa. Their functions are not defined, but they may protect the mucosa from insults, stabilize the mucus layer, and




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




affect healing of the epithelium. This gene, which is expressed in the gastric mucosa, has also been studied because of its expression in human tumors. This gene and two other related trefoil family member genes are found in a cluster on chromosome 21. [provided by RefSeq, Jul 2008],


Immunohistochemical analysis of paraffin-embedded human-breast-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-breast-cancer, antibody was diluted at 1:200




Immunohistochemical analysis of paraffin-embedded human-stomach, antibody was diluted at 1:200



+86-27-59760950







Immunohistochemical analysis of paraffin-embedded human-stomach, antibody was diluted at 1:200



+86-27-59760950