

## Frizzled-4 rabbit pAb

Cat No.:ES8800

For research use only

## Overview

Product Name Frizzled-4 rabbit pAb

Host species Rabbit
Applications IHC;IF;ELISA

Species Cross-Reactivity Human; Mouse; Rat

Recommended dilutions IHC-p 1:50-200, ELISA 1:10000-20000

**Immunogen** Synthetic peptide from human protein at AA range:

11-60

**Specificity** The antibody detects endogenous Frizzled-4

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Frizzled-4 (Fz-4) (hFz4) (FzE4) (CD antigen CD344)

Gene Name FZD4

Cellular localizationCell membrane ; Multi-pass membrane protein .PurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 8322 Human Swiss-Prot Number Q9ULV1

Alternative Names Frizzled-4 (Fz-4;hFz4;FzE4;CD antigen CD344)

Background frizzled class receptor 4(FZD4) Homo sapiens

This gene is a member of the frizzled gene family.

Members of this family encode


seven-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled

receptors are coupled to the beta-catenin canonical signaling pathway. This protein may play a role as a positive regulator of the Wingless type MMTV integration site signaling pathway. A transcript




variant retaining intronic sequence and encoding a shorter isoform has been described, however, its expression is not supported by other experimental evidence. [provided by RefSeq, Jul 2008],

Immunohistochemical analysis of paraffin-embedded Human-skin, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-skin, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-brain, antibody was diluted at 1:100

