

## IL-17F rabbit pAb

## Cat No.:ES8805

For research use only

## Overview

Product Name	IL-17F rabbit pAb
Host species	Rabbit
Applications	IHC;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	IHC-p 1:50-200, ELISA 1:10000-20000
Immunogen	Synthetic peptide from human protein at AA range: 81-130
Specificity	The antibody detects endogenous IL-17F
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Interleukin-17F (IL-17F) (Cytokine ML-1)
	(Interleukin-24) (IL-24)
Gene Name	IL17F IL24
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	
Human Gene ID	112744
Human Swiss-Prot Number	Q96PD4
Alternative Names	Interleukin-17F (IL-17F;Cytokine
	ML-1;Interleukin-24;IL-24)
Background	The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM_CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte



chemoattractant protein-1. [provided by RefSeq, Jul 2008],

Immunohistochemical analysis of paraffin-embedded Human-colon-cancer, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-colon-cancer, antibody was diluted at 1:100

