

## **INCE** rabbit pAb

**Cat No.:ES9002** 

For research use only

## Overview

Product Name INCE rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein . at

AA range: 1-80

**Specificity** INCE Polyclonal Antibody detects endogenous levels

of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at  $-20^{\circ}$ C. Avoid repeated freeze-thaw cycles.

Protein Name Inner centromere protein

Gene Name INCENP

Cellular localization Nucleus . Chromosome, centromere . Cytoplasm,

cytoskeleton, spindle. Midbody. Chromosome,

centromere, kinetochore. Colocalized at synaptonemal complex central element from zygotene up to late pachytene when it begins to relocalize to heterochromatic chromocenters. Colocalizes with AURKB at a connecting strand traversing the centromere region and joining sister kinetochores, in metaphase II centromeres. This strand disappears at the metaphase II/anaphase II transition and relocalizes to the spindle midzone (By similarity). Colocalizes with AURKB at mitotic

similarity). Colocalizes with AURKB at mitotic chromosomes (PubMed:11453556). Localizes to inner kinetochore (PubMed:16760428). Localizes on chromosome arms and inner centromeres from prophase through metaphase and then transferring to the spindle midzone and midbody from anaphase

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using





+86-27-59760950 ELk

ELKbio@ELKbiotech.com

www.elkbiotech.com



epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 100kD
Human Gene ID 3619
Human Swiss-Prot Number Q9NQS7

**Alternative Names** 

**Background** 

In mammalian cells, 2 broad groups of centromere-interacting proteins have been described: constitutively binding centromere proteins and 'passenger,' or transiently interacting, proteins (reviewed by Choo, 1997). The constitutive proteins include CENPA (centromere protein A; MIM 117139), CENPB (MIM 117140), CENPC1 (MIM 117141), and CENPD (MIM 117142). The term 'passenger proteins' encompasses a broad collection of proteins that localize to the centromere during specific stages of the cell cycle (Earnshaw and Mackay, 1994 [PubMed 8088460]). These include CENPE (MIM 117143); MCAK (MIM 604538); KID (MIM 603213); cytoplasmic dynein (e.g., MIM 600112); CliPs (e.g., MIM 179838); and CENPF/mitosin (MIM 600236). The inner centromere proteins (INCENPs) (Earnshaw and Cooke, 1991

[PubMed 1860899]), the initial members of the passenger protein group, display a broad localization



alo