



# CDA7L rabbit pAb

Cat No.:ES9119

For research use only

## Overview

<b>Product Name</b>	CDA7L rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;ELISA
<b>Species Cross-Reactivity</b>	Human;Rat;Mouse;
<b>Recommended dilutions</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 140-220
<b>Specificity</b>	CDA7L Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C . Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	Cell division cycle-associated 7-like protein (Protein JPO2) (Transcription factor RAM2)
<b>Gene Name</b>	CDCA7L HR1 JPO2 R1
<b>Cellular localization</b>	Cytoplasm. Nucleus. Associates with chromatin. Translocates from cytoplasm to nucleus under dexamethasone induction.
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	49kD
<b>Human Gene ID</b>	55536
<b>Human Swiss-Prot Number</b>	Q96GN5
<b>Alternative Names</b>	
<b>Background</b>	function:Plays a role in transcriptional regulation as a repressor that inhibits monoamine oxidase A (MAOA) activity and gene expression by binding to the promoter. Plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. Involved in apoptotic





**ELK Biotechnology**

signaling pathways; May act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1.,induction:By MYC overexpression in a concentration dependent manner in neuroblastoma cell line.,miscellaneous:Cells lacking CDCA7L display a reduction of 25-30% of colony formation in medulloblastoma cell lines. CDCA7L overexpression induces colony formation.,subcellular location:Associates with chromatin. Translocates from cytoplasm to nucleus under dexamethasone induction.,subunit:Interacts with MYC and PSIP1.,tissue specificity:Ubiquitous. Overexpressed in medulloblastoma.,



+86-27-59760950

[ELKbio@ELKbiotech.com](mailto:ELKbio@ELKbiotech.com)

[www.elkbiotech.com](http://www.elkbiotech.com)

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C