

MED10 rabbit pAb

Cat No.: ES9785

For research use only

Overview

Product Name MED10 rabbit pAb

Host species Rabbit
Applications WB;ELISA
Species Cross-Reactivity Human;Mouse

Recommended dilutions WB 1:500-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human protein .

at AA range: 40-120

Specificity MED10 Polyclonal Antibody detects endogenous

levels of protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Mediator of RNA polymerase II transcription subunit

10 (Mediator complex subunit 10)

(Transformation-related gene 17 protein) (TRG-17) (Transformation-related gene 20 protein) (TRG-20)

Gene Name MED10 L6 TRG17 TRG20

Cellular localization Nucleus.

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band14kDHuman Gene ID84246Human Swiss-Prot NumberQ9BTT4

Alternative Names

Background MED10 is a component of the Mediator complex,

which is a coactivator for DNA-binding factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed 12584197]).[supplied by OMIM,


Oct 2008],



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

