



NBAS rabbit pAb

Cat No.:ES9903

For research use only

Overview

| | |
|---------------------------------|---|
| Product Name | NBAS rabbit pAb |
| Host species | Rabbit |
| Applications | IHC;IF |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | IHC-p 1:50-300 |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 1880-1960 |
| Specificity | NBAS Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Neuroblastoma-amplified sequence (Neuroblastoma-amplified gene protein) |
| Gene Name | NBAS NAG |
| Cellular localization | Cytoplasm . Endoplasmic reticulum . Endoplasmic reticulum membrane; Peripheral membrane protein . |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | 260kD |
| Human Gene ID | 51594 |
| Human Swiss-Prot Number | A2RRP1 |
| Alternative Names | |
| Background | This gene encodes a protein with two leucine zipper domains, a ribosomal protein S14 signature domain and a Sec39 like domain. The protein is thought to be involved in Golgi-to-ER transport. Mutations in this gene are associated with short stature, optic nerve atrophy, and Pelger-Huet anomaly. [provided |






ELK Biotechnology

by RefSeq, Oct 2012],

Immunohistochemical analysis of paraffin-embedded human small intestinal carcinoma tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:2



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C