

GDNF rabbit pAb

Cat No.:ES11350

For research use only

Overview

Product Name	GDNF rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB 1:500-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human protein .
	at AA range: 100-180
Specificity	GDNF Polyclonal Antibody detects endogenous
	levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Glial cell line-derived neurotrophic factor (hGDNF)
	(Astrocyte-derived trophic factor) (ATF)
Gene Name	GDNF
Cellular localization	Secreted .
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	23kD
Human Gene ID	2668
Human Swiss-Prot Number	P39905
Alternative Names	
Background	This gene encodes a secreted ligand of the TGF-beta
	(transforming growth factor-beta) superfamily of
	proteins. Ligands of this family bind various
	TGF-beta receptors leading to recruitment and
	activation of SMAD family transcription factors that
	regulate gene expression. The encoded
	preproprotein is proteolytically processed to
	generate each subunit of the disulfide-linked



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



homodimer. The recombinant form of this protein, a highly conserved neurotrophic factor, was shown to promote the survival and differentiation of dopaminergic neurons in culture, and was able to prevent apoptosis of motor neurons induced by axotomy. This protein is a ligand for the product of the RET (rearranged during transfection) protooncogene. Mutations in this gene may be associated with Hirschsprung disease and Tourette syndrome. This gene encodes multiple protein isoforms that may undergo similar proteolytic

Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C