

O11H1 rabbit pAb

Cat No.:ES14414

For research use only

Overview

O11H1 rabbit pAb
Rabbit
WB
Human; Rat; Mouse;
WB 1: 500-2000
Synthesized peptide derived from human O11H1 AA range: 187-237
This antibody detects endogenous levels of O11H1 at Human
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Store at -20°C. Avoid repeated freeze-thaw cycles.
O11H1
OR11H1
Cell membrane; Multi-pass membrane protein.
The antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using
epitope-specific immunogen.
Polyclonal
1 mg/ml
81061
Q8NG94
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C