

CD11A rabbit pAb

Cat No.:ES17739

For research use only

Overview

| | |
|--------------------------|---|
| Product Name | CD11A rabbit pAb |
| Host species | Rabbit |
| Applications | WB |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | WB 1: 500-2000 |
| Immunogen | Synthesized peptide derived from human CD11A AA range: 53-103 |
| Specificity | This antibody detects endogenous levels of CD11A at Human |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | CD11A |
| Gene Name | CDK11A CDC2L2 CDC2L3 PITSLRB |
| Cellular localization | Cytoplasm. Nucleus. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | |
| Human Gene ID | 728642 |
| Human Swiss-Prot Number | Q9UQ88 |
| Alternative Names | |
| Background | This gene encodes a member of the serine/threonine protein kinase family. Members of this kinase family are known to be essential for eukaryotic cell cycle control. Due to a segmental duplication, this gene shares very high sequence identity with a neighboring gene. These two genes are frequently deleted or altered in neuroblastoma. The protein kinase encoded by this gene can be cleaved by caspases and may play a role in cell |



apoptosis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015],

Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night

