

## Caspase 10 (p12, Cleaved-Ala416) rabbit pAb

Cat No.: ES19952

For research use only

Overview

Product Name Caspase 10 (p12, Cleaved-Ala416) rabbit pAb

Host species Rabbit
Applications WB; ELISA

**Species Cross-Reactivity** Human;Rat;Mouse;

Recommended dilutions WB 1:1000-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from human Caspase

10 (p12, Cleaved-Ala416)

**Specificity** This antibody detects endogenous levels of Human

Caspase 10 (p12, Cleaved-Ala416, protein was cleaved amino acid sequence between 415-416)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at  $-20^{\circ}$ C. Avoid repeated freeze-thaw cycles.

Protein Name Caspase10
Gene Name CASP10 MCH4

**Cellular localization** 

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 12 55kD
Human Gene ID 843
Human Swiss-Prot Number Q92851

Alternative Names Caspase-10 (CASP-10;EC 3.4.22.63;Apoptotic

protease Mch-4;FAS-associated death domain protein interleukin-1B-converting enzyme

2;FLICE2;ICE-like apoptotic protease 4) [Cleaved into: Caspase-10 subunit p23/17; Caspase-10 subunit

p12]

**Background** This gene encodes a protein which is a member of

the cysteine-aspartic acid protease (caspase) family.

+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with type IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Apr 2011],



+86-27-59760950