

## Caspase-10 p23/17 (Cleaved-D415) rabbit

## pAb

Cat No.:ES19953

For research use only

## Overview

Product Name	Caspase-10 p23/17 (Cleaved-D415) rabbit pAb
Host species	Rabbit
Applications	WB; ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
<b>Recommended dilutions</b>	WB 1:1000-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human
	Caspase-10 p23/17 (Cleaved-D415)
Specificity	This antibody detects endogenous levels of Human
	Caspase-10 p23/17 (Cleaved-D415, protein was
	cleaved amino acid sequence between 415-416)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20 $^\circ\!\mathrm{C}$ . Avoid repeated freeze-thaw cycles.
Protein Name	Caspase10
Gene Name	CASP10 MCH4
Cellular localization	
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	23/17 55kD
Human Gene ID	843
Human Swiss-Prot Number	Q92851
Alternative Names	Caspase-10 (CASP-10;EC 3.4.22.63;Apoptotic
	protease Mch-4;FAS-associated death domain
	protein interleukin-1B-converting enzyme
	2;FLICE2;ICE-like apoptotic protease 4) [Cleaved into:
	Caspase-10 subunit p23/17; Caspase-10 subunit
	p12]
Background	This gene encodes a protein which is a member of
-	the cysteine-aspartic acid protease (caspase) family.
+86-27-59760950	ELKbio@ELKbiotech.com www.elkbiotech.com



23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with type IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Apr 2011],



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C