

## Histone H4 (Phospho Thr80) rabbit pAb

## Cat No.:ES20564

For research use only

## Overview

Product Name	Histone H4 (Phospho Thr80) rabbit pAb
Host species	Rabbit
Applications	WB
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	WB: 1:1000-2000
Immunogen	Synthetic Peptide of Histone H4 (Phospho Thr80)
Specificity	The antibody detects endogenous Histone H4
	(Phospho Thr80) protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Histone H4
Gene Name	HIST1H4A
Cellular localization	Nucleus. Chromosome.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	
Observed band	14kD
Human Gene ID	121504
Human Swiss-Prot Number	P62805
Alternative Names	HIST1H4A; H4/A; H4FA; HIST1H4B; H4/I; H4FI;
	HIST1H4C; H4/G; H4FG; HIST1H4D; H4/B; H4FB;
	HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC;
	HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM;
	HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD;
	HIST1H4L; H4/K; H4FK; HIST2H4A; H4/
Background	Histones are basic nuclear proteins that are
	responsible for the nucleosome structure of the
	chromosomal fiber in eukaryotes. Two molecules of
	each of the four core histones (H2A, H2B, H3, and
	H4) form an octamer, around which approximately



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015],

Western blot analysis of extracts from Hela cells, untreated (-) or treated, 1:5000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C