

## Olfactory receptor 52B2 rabbit pAb

Cat No.: ES3052

For research use only

## Overview

Product Name Olfactory receptor 52B2 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

**Species Cross-Reactivity** Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/5000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR52B2. AA

range:201-250

**Specificity** Olfactory receptor 52B2 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 52B2

protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 52B2

Gene Name OR52B2

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit<br/>antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band36kDHuman Gene ID255725Human Swiss-Prot NumberQ96RD2

Alternative Names OR52B2; Olfactory receptor 52B2; Olfactory

receptor OR11-70

**Background** Olfactory receptors interact with odorant molecules

in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of




+86-27-59760950 ELKbio@ELKbiotech.com www.elk




G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Olfactory receptor 52B2 Polyclonal Antibody




Immunofluorescence analysis of MCF7 cells, using OR52B2 Antibody. The picture on the right is blocked with the synthesized peptide.




+86-27-59760950







Western blot analysis of lysates from COLO, HUVEC, and HepG2 cells, using OR52B2 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

Western blot analysis of the lysates from HepG2 cells using OR52B2 antibody.

