

Caspase-10 rabbit pAb

Cat No.:ES3824

For research use only

Overview

Product Name	Caspase-10 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not
	yet tested in other applications.
Immunogen	Synthesized peptide derived from the Internal
_	region of human Caspase-10.
Specificity	Caspase-10 Polyclonal Antibody detects endogenous
	levels of Caspase-10 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Caspase10
Gene Name	CASP10
Cellular localization	cytosol,CD95 death-inducing signaling
	complex,ripoptosome,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	58kD
Human Gene ID	843
Human Swiss-Prot Number	Q92851
Alternative Names	CASP10; MCH4; Caspase-10; CASP-10; Apoptotic
	protease Mch-4; FAS-associated death domain
	protein interleukin-1B-converting enzyme 2; FLICE2;
	ICE-like apoptotic protease 4
Background	This gene encodes a protein which is a member of
	the cysteine-aspartic acid protease (caspase) family.
	Sequential activation of caspases plays a central role
	in the execution-phase of cell apoptosis. Caspases



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with type IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Apr 2011],

Western Blot analysis of extracts from Jurkat, K562 cells, using Caspase-10 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C