

β-1,4-Gal-T2 rabbit pAb

Cat No.:ES7862

For research use only

Overview

| Product Name | β-1,4-Gal-T2 rabbit pAb |
|------------------------------|---|
| Host species | Rabbit |
| Applications | IHC;IF;ELISA |
| Species Cross-Reactivity | Human;Mouse |
| Recommended dilutions | Immunohistochemistry: 1/100 - 1/300. ELISA: |
| | 1/20000. Not yet tested in other applications. |
| Immunogen | Synthesized peptide derived from the C-terminal |
| | region of human β-1,4-Gal-T2. |
| Specificity | β-1,4-Gal-T2 Polyclonal Antibody detects |
| | endogenous levels of β-1,4-Gal-T2 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and |
| | 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | Beta-1,4-galactosyltransferase 2 |
| Gene Name | B4GALT2 |
| Cellular localization | Golgi apparatus, Golgi stack membrane; Single-pass |
| | type II membrane protein. Trans cisternae of Golgi |
| | stack. |
| Purification | The antibody was affinity-purified from rabbit |
| | antiserum by affinity-chromatography using |
| | epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | |
| Human Gene ID | 8704 |
| Human Swiss-Prot Number | O60909 |
| Alternative Names | B4GALT2; Beta-1; 4-galactosyltransferase 2; |
| | Beta-1,4-GalTase 2; Beta4Gal-T2; b4Gal-T2; |
| | UDP-Gal:beta-GlcNAc beta-1,4-galactosyltransferase |
| | 2; UDP-galactose:beta-N-acetylglucosamine |
| | beta-1,4-galactosyltransferase 2 |
| Background | This gene is one of seven |
| | beta-1,4-galactosyltransferase (beta4GalT) genes. |



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. The enzyme encoded by this gene synthesizes N-acetyllactosamine in glycolipids and glycoproteins. Its substrate specificity i

Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C