

# Histone H3 (Di-Methyl-K80) rabbit pAb

Cat No.:ES8831

For research use only

## Overview

|                              |  |
|------------------------------|--|
| <b>Product Name</b>          | Histone H3 (Di-Methyl-K80) rabbit pAb  |
| <b>Host species</b>          | Rabbit   |
| <b>Applications</b>          | WB;ELISA   |
| <b>Species</b>               | Human:K80Mouse:K80Rat:K80  |
| <b>Cross-Reactivity</b>      |  |
| <b>Recommended dilution</b>  | wb dilution 1:1000   |
| <b>Immunogen</b>             | Synthesized Tri-Methyl peptide derived from human Histone H3. at AA range: K80   |
| <b>Specificity</b>           | This antibody detects endogenous levels of Histone H3 at Human:K80Mouse:K80Rat:K80, It doesn't react with total protein.         |
| <b>Formulation</b>           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| <b>Storage</b>               | Store at -20°C. Avoid repeated freeze-thaw cycles.   |
| <b>Protein Name</b>          | Histone H3   |
| <b>Gene Name</b>             | HIST1H3A/HIST1H3B/HIST1H3C/HIST1H3D/HIST1H3E/HIST1H3F/HIST1H3G/HIST1H3H/HIST1H3I/HIST1H3J/HIST2H3A/HIST2H3C/HIST2H3D/H3F3A/H3F3B |
| <b>Cellular localization</b> | Nucleus. Chromosome.   |
| <b>Purification</b>          | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.            |
| <b>Clonality</b>             | Polyclonal   |





Y

**Concentration** 1 mg/ml

**Observed band**

**Human Gene ID** 8350

**Human Swiss-Prot Number**


P68431/Q71DI3/P84243/Q6NXT2

**Alternative Names**

H3K80ME2; HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3/a; Histone H3/b; Histone H3/c; Histone H3/d; Histone H3/f; H

**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015],



Western Blot analysis of HeLa cells using primary antibody diluted at 1:1000 (4°C overnight). Secondary antibody: Goat Anti-rabbit IgG IRDye 800 (diluted at 1:5000, 25°C, 1 hour)

