

## **BCL3** rabbit pAb

Cat No.: ES9000

For research use only

## Overview

**Product Name** BCL3 rabbit pAb

**Host species** Rabbit IHC;IF;WB **Applications** 

**Species Cross-Reactivity** Human;Rat;Mouse;;Goat

**Recommended dilutions** WB 1:500-2000 ELISA 1:5000-20000

**Immunogen** Synthesized peptide derived from human protein . at

AA range: 340-420

BCL3 Polyclonal Antibody detects endogenous levels Specificity

of protein.

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at  $-20^{\circ}$ C. Avoid repeated freeze-thaw cycles. **Storage** B-cell lymphoma 3 protein (BCL-3) (Proto-oncogene **Protein Name** 

BCL3)

**Gene Name** BCL3 BCL4 D19S37

Cellular localization Nucleus. Cytoplasm. Cytoplasm, perinuclear region.

Ubiquitination via 'Lys-63'-linked ubiquitin chains is

required for nuclear accumulation. .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 49kD **Human Gene ID** 602 **Human Swiss-Prot Number** P20749

**Alternative Names** 

Background This gene is a proto-oncogene candidate. It is

identified by its translocation into the

immunoglobulin alpha-locus in some cases of B-cell leukemia. The protein encoded by this gene contains

seven ankyrin repeats, which are most closely related to those found in I kappa B proteins. This



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



protein functions as a transcriptional co-activator that activates through its association with NF-kappa B homodimers. The expression of this gene can be induced by NF-kappa B, which forms a part of the autoregulatory loop that controls the nuclear residence of p50 NF-kappa B. [provided by RefSeq, Jul 2008],



+86-27-59760950